

Factors Associated with Clinical Outcomes in Spontaneous Subarachnoid Hemorrhage

Spontan Subaraknoid Kanamada Klinik Sonuçlarla İlişkili Faktörler

Ömer Özdemir, Furkan Diren, Osman Boyalı, Benan Baysoy, Serdar Kabataş

University of Health Sciences Turkey, Gaziosmanpaşa Training and Research Hospital, Clinic of Neurosurgery, Istanbul, Turkey

Cite this article as: Özdemir Ö, Diren F, Boyalı O, Baysoy B, Kabataş S. Factors Associated with Clinical Outcomes in Spontaneous Subarachnoid Hemorrhage. J Acad Res Med 2022;12(3):124-9

ABSTRACT

Objective: This study aims at evaluating the factors affecting prognosis in patients with aneurysmal subarachnoid hemorrhage (SAH).

Methods: The charts and 6th month outpatient clinic records of the patients who were treated with the diagnosis of SAH between 2016 and 2020 were retrospectively reviewed. Post-discharge clinical status, adaptation to life, and disability status of 44 patients with aneurysmal SAH who were treated with endovascular or microsurgical techniques in our clinic were evaluated with the Glasgow Outcome Scale Extended (GOSE). The relationship between the patient's age, gender, timing of treatment, the location of the aneurysm, the World Federation of Neurological Surgeons (WFNS) scale score, Hunt-Hess scale (HHS) score, Fisher scale score and preoperative evaluations that might affect the GOSE score after discharge were examined.

Results: Forty-four patients who met the criteria were included in the study. The mean age of the patients was 49.1±10.3 (minimum-maximum: 29-77). Of those, 54.5% were females while 45.5% were males. Comorbidity was present in 54.5% of the patients. The most common aneurysm was the middle cerebral artery aneurysm with the rate of 27.3%. Of the aneurysmal SAHs, 45.5% were treated with microsurgical clipping and 27.3% with endovascular treatment. A statistically significant correlation was found between GOSE score and timing of treatment days (p=0.014), WFNS (p=0.002) and HHS scores (p<0.001). No statistically significant correlation was found between GOSE score and patient's age (p=0.47) and Fisher scale score (p=0.465).

Conclusion: In our study, the effects of early surgery, WFNS, and HHS scores on clinical outcome in cerebral aneurysmatic SAH were shown.

Keywords: Spontaneous subarachnoid hematoma, prognosis, clinical outcomes

ÖZ

Amaç: Bu çalışmada, anevrizmal subaraknoid kanamalı (SAK) hastalarda prognozu etkileyen faktörlerin değerlendirilmesi amaçlanmaktadır.

Yöntemler: Bu çalışmada 2016-2020 yılları arasında SAK tanısı ile tedavi edilen hastaların hasta dosyaları ve taburculuk sonrası 6. ay poliklinik kayıtları geriye dönük olarak incelendi. Kliniğimizde endovasküler veya mikrocerrahi tekniklerle tedavi edilen 44 anevrizmal SAK'li hastanın taburculuk sonrası klinik durumu, hayata adaptasyonu ve maluliyet durumu Glasgow Outcome Scale-Extended (GOSE) ile değerlendirildi. Hastanın yaşı, cinsiyeti, tedavi zamanlaması, anevrizmanın yeri, Dünya Nörolojik Cerrahlar Federasyonu (WFNS) skalası, Hunt-Hess skalası (HHS), Fisher skalası ve sonrasında GOSE skorunu etkileyebilecek preoperatif değerlendirmeler arasındaki ilişki incelendi.

Bulgular: Kriterleri karşılayan 44 hasta çalışmaya dahil edildi. Hastaların yaş ortalaması 49,1±10,3 (minimum-maksimum: 29-77) idi. Bunların %54,5'i kadın, %45,5'i erkekti. Hastaların %54,5'inde komorbidite mevcuttu. En sık görülen anevrizma %27,3 oranı ile orta serebral arter anevrizması idi. Anevrizmal SAH'lerin %45,5'i mikrocerrahi klipsleme ile ve %27,3'ü endovasküler yolla tedavi edildi. GOSE skoru ile tedavinin zamanlaması (p=0,014), WFNS (p=0,002) ve HHS skorları (p<0,001) arasında istatistiksel olarak anlamlı bir ilişki bulunurken, hasta yaşı (p=0,47) ve Fisher skalası skoru (p=0,465) ile istatistiksel anlamlı ilişki bulunmadı.

Sonuç: Çalışmamızda serebral anevrizmatik SAK'de erken cerrahi günü, WFNS ve HHS'nin klinik sonuca etkileri gösterilmiştir.

Anahtar kelimeler: Spontan subaraknoid hematoma, prognoz, klinik sonuçlar

ORCID IDs of the authors: Ö.Ö. 0000-0003-3783-0203; F.D. 0000-0001-6169-9722; O.B. 0000-0002-2500-1718; B.B. 0000-0001-9536-3768; S.K. 0000-0003-2691-6861.

Corresponding Author/Sorumlu Yazar: Ömer Özdemir,

E-mail: dromerozdemir1982@gmail.com



Received Date/Geliş Tarihi: 13.06.2022 **Accepted Date/Kabul Tarihi:** 04.10.2022

©Copyright 2022 by University of Health Sciences Turkey, Gaziosmanpaşa Training and Research Hospital. Available on-line at www.jarem.org

©Telif Hakkı 2022 Sağlık Bilimleri Üniversitesi, Gaziosmanpaşa Eğitim ve Araştırma Hastanesi. Makale metnine www.jarem.org web sayfasından ulaşılabilir.

INTRODUCTION

Cerebral aneurysms (CA) occur in 3-5% of the general population (1). Aneurysm rupture causes subarachnoid hemorrhage (SAH) and is associated with high mortality and morbidity (2). Its formation and pathophysiology have not been fully defined (3). Current treatment options consist of endovascular treatment and microsurgical clipping. Despite advances in surgical and endovascular techniques, equipment diversity in devices, and the improvements in intensive care units, mortality and morbidity associated with aneurysm rupture remain high (4). Treating ruptured aneurysms is required. The type of treatment is decided according to the choice of patients, the size and site of aneurysm. However, at vascular centers with a high annual number of microsurgical clipping, there is a tendency to perform clipping rather than applying endovascular treatment (5). Although the results of the International Study of Unruptured Intracranial Aneurysms and International Subarachnoid Aneurysm studies point to "coil first" with endovascular surgical intervention, there are ongoing discussions regarding the superiority of endovascular treatment and microsurgical clipping treatment over each other and which one should be preferred (6,7).

The Glasgow outcome scale-extended (GOSE) is used in the treatment of SAH and in the short and long-term follow-ups and results of these treatments, as in other cranial pathologies with brain injury (8,9). Many studies in the literature evaluate clinical outcome with GOSE. Correlation of applied treatment parameters and neurological status with GOSE is examined (8,9). Recently, some studies claimed that early aneurysm treatment achieves better clinical outcomes (8,10). There are also studies showing poor clinical outcomes in patients with WFNS grade IV and V SAH (9). There are few studies comparing GOSE score with preoperative parameters in SAH. No study was found in which all the mentioned parameters were compared with GOSE score.

In this study, we planned to investigate the effects of preoperative parameters and treatments in spontaneous SAH on clinical outcome.

METHODS

The data of 44 patients who developed SAH due to CA rupture and were admitted to our clinic and were treated between 2016-2020 were analysed retrospectively. The patients with spontaneous SAH who were admitted to our hospital within 72 hours after bleeding were included in the study. The patients with non-spontaneous SAH and those with previous record of neurological diseases (cerebrovascular disease, neurodegenerative diseases, etc.) were excluded from the study. All data of the patients were collected from the hospital database. Data were collected in the following categories: patient age, gender, World Federation of Neurological Surgeons (WFNS) scale score (Table 1) (11), Hunt-Hess scale (HHS) score (Table 2) (12), Fisher scale score (Table 3) (12), the site of aneurysm, the type of treatment performed (microsurgical clipping/endovascular treatment) and the number of days between first bleeding and application of treatment. In

addition, the patients' adaptation to life and disability status were evaluated with the (GOSE) score (Table 4) (13) at the 6th month after discharge. Cerebral computed tomography (CT), cerebral CT angiography and Digital Subtraction Angiography (DSA) were performed in all patients after admission to the hospital. Control DSA was also performed in the patients at the 6th month.

The patients were treated by a single surgeon experienced in vascular surgery. They were operated with micro-Doppler and neuromonitoring.

Single or multiple aneurysm clips (Sugita®, Mizuho, Tokyo, Japan) were used during surgery. Endovascular treatment was performed by experienced senior neuroradiologists. Diagnostic or therapeutic angiography was performed with a routine femoral approach using biplane angio-graphic system with a flat panel detector (Toshiba Infinix, Toshiba Medical, Nasu, Japan). Lesions were treated with coils (Boston Scientific Corp., Natick, MA, USA) and stent (Neuroform® Stent System, Boston Scientific Corp., Natick, MA, USA).

All procedures were carried out in accordance with the 1964 Helsinki declaration. Written informed consent for scientific

Table 1. World Federation of Neurological Surgeons scale

Grade	Glasgow coma scale score	Motor deficit
I	15	Absent
II	13-14	Absent
III	13-14	Present
IV	7-12	Present or absent
V	3-6	Present or absent

Table 2. Hunt-Hess scale

Grade	Criteria
I	Asymptomatic, or minimal headache, nuchal rigidity
II	Moderate to severe headache, no neurological deficit except for cranial nerve palsy
II	Drowsiness, confusion, mild focal deficit
IV	Stupor, moderate to severe hemiparesis, early decerebrate posturing
V	Deep coma, decerebrate posturing, moribund

Table 3. Fisher scale

Grade	CT scan
1	No blood visualized
2	A diffuse deposition or thin layer with all vertical layers of blood (interhemispheric fissure, insular cistern, ambient cistern) less than 1 mm thick
3	Localized clots and/or vertical layers of blood 1 mm or greater in thickness
4	Diffuse or no subarachnoid blood, but with intracerebral or intraventricular clots

CT: computed tomography

Table 4. Glasgow outcome scale-extended

Category	Name	Definition
1	Death	
2	Persistent vegetative state	Unresponsive and speechless
3	Sever disability Lower	Requires frequent help of someone to be around at home most of the time every day
4	Sever disability Upper	Can be left alone >8 h during the day, but unable to travel and/or go shopping without assistance
5	Moderate disability Lower	Unable to work or only in sheltered workshop
6	Moderate disability Upper	Reduced work capacity; resumes <50% of the pre-injury level of social and leisure activities
7	Good recovery Lower	Minor problems that affect daily life; resumes >50% of the pre-injury level of social and leisure activities
8	Good recovery Upper	No current problems related to the brain injury that affect daily life

Table 5. Description of study participants

	Characteristics	
Age	$\bar{x}\pm s$	49.1±10.3
Gender	Female, n (%)	24 (54.5)
	Male, n (%)	20 (45.5)
Co-morbidities	Yes, n (%)	24 (54.5)
	No, n (%)	20 (45.5)
Hypertension	Yes, n (%)	20 (45.5)
	No, n (%)	24 (54.5)
Asthma	Yes, n (%)	4 (9.1)
	No, n (%)	40 (90.9)
Other co-morbidities	Yes, n (%)	10 (22.7)
	No, n (%)	34 (77.3)

Table 6. WFNS scale, HHS and Fisher scale scores

	Score	n (%)
WFNS scale	1	18 (40.9)
	2	10 (22.7)
	3	8 (18.2)
	4	5 (11.4)
	5	3 (6.8)
Fisher scale	1	3 (6.8)
	2	8 (18.2)
	3	17 (38.6)
	4	16 (36.4)
HHS	1	18 (40.9)
	2	10 (22.7)
	3	8 (18.2)
	4	5 (11.4)
	5	3 (6.8)

WFNS: World Federation of Neurological Surgeons, HHS: Hunt-Hess scale

purposes and clinical data collection were obtained from patients according to institutional protocol. Ethical approval was obtained from the local ethics committee (approval no: 319, date: 22.09.2021).

Statistical Analysis

The Statistical Package for Social Sciences (SPSS) 25.0 was used for statistical analysis (SPSS Inc.; Chicago, IL, USA). The Kolmogorov-Smirnov test was used to examine the normal distribution. According to results of normality analyses, the data was not normally distributed. The descriptive statistical methods (frequency, percentage, mean, standard deviation) were used to evaluate the demographic data. Pearson's chi-square test was used to compare the qualitative data. The Spearman correlation analysis was used for analysing the association of the quantitative data. The results were evaluated at a confidence interval of 95% and a significance level of $p<0.05$.

RESULTS

The total number of patients who were admitted to the clinic with the diagnosis of CA and underwent treatment between 2016 and 2020 was 44. The characteristic features of the patients are given in Table 5. Their mean age was 49.1±10.3 (minimum-maximum: 29-77). Of them 54.5% were females while 45.5% were males. Comorbidity was present in 54.5% of the patients. The three most common comorbidities were hypertension (HT) (45.5%), asthma and chronic obstructive pulmonary disease. WFNS scale, HHS and Fisher scale scores of the patients are shown in Table 6. The most common values for each scale were as follows: For WFNS scale score "Grade 1" in 18 (40.9%) patients, for HHS score "1" in 18 (40.9%) patients, and for Fisher score "3" in 17 (38.6%) patients.

The angiography results of the patients with SAH were as follows: Twelve (27.3%) had middle cerebral artery (MCA) aneurysm, 8 (18.2%) anterior communicating artery (AcoA) aneurysm, 4 (9.1%) internal carotid artery aneurysm, 3 (6.8%) posterior communicating artery aneurysm, 2 (4.5%) distal anterior cerebral artery aneurysm, 2 (4.5%) posterior inferior cerebellar artery aneurysm, 1 (2.3%)

basilar artery type aneurysm (Figure 1). Twelve (27.3%) patients were angiography negative. Except for these 12 patients, 20 (45.5%) patients were treated with microsurgical clipping and 12 (27.3%) patients with endovascular treatment, making the total number of treated patients 32. The mean number of days between bleeding and application of microsurgical clipping or endovascular treatment after SAH was 1.2, with the first day and the fifth day of bleeding representing the minimum and maximum values, respectively.

The study found a statistically significant negative correlation between GOSE score and timing of treatment ($p=0.014$) and a moderate significant negative correlation between GOSE and WFNS scale ($p=0.002$) and HHS scores ($p<0.001$) (Table 7). The overall mortality rate of the patients at the 6-month follow-up was 11.3%.

DISCUSSION

Nearly half of the SAH cases are caused by CA (14). Studies on predicting the prognosis of patients after SAH and the risk factors that affect the treatment outcome are still pending (15). In our current study, a negative correlation was found between WFNS scale and HHS scores, which were the preoperative evaluation scales, the GOSE score, and the timing of treatment applied.

Table 7. Relationship of clinical parameters with GOSE

	GOSE	
	r	p-value*
Surgery day	-0.390	0.014
WFNS	-0.477	0.002
HHS	-0.566	<0.001
Age	0.119	0.47
Fisher scale	-0.120	0.465

*Spearman correlation, GOSE: Glasgow outcome scale-extended, WFNS: World Federation of Neurological Surgeons, HHS: Hunt-Hess scale

ANGIOGRAPHY RESULTS (PERCENTILE)

■ Angio (-) ■ MCA ■ AcoA ■ ICA ■ PcoA ■ ACA ■ PICA ■ Basilar tip




Figure 1. Angiography results (percentile)

MCA: middle cerebral artery, AcoA: anterior communicating artery, ICA: internal carotid artery, PcoA: posterior communicating artery, ACA: anterior cerebral artery, PICA: posterior inferior cerebellar artery

In this study, 27.3% of the 44 patients with SAH were found to have negative angiogram. When compared with the literature this rate is high. In the literature, angiogram negative SAHs constitute approximately 10-20% of spontaneous SAHs (16,17). It also has the same negative clinical outcomes as aneurysmal SAH (16,17). In line with the literature, GOSE scores of patients with angiogram negative SAH and aneurysmal SAH are similar. It is thought that the frequency of angiogram negative SAHs has increased, and this may be related to the widespread use of antiplatelet drugs (18).

Complications after SAH (vasospasm, ischemic neurological deficit, and brain infarction) are closely related to age and gender (19). While the incidence of complications is lower in elderly patients, they occur more commonly among the younger patients (20). It has been stated that 30-39 age group has higher risk whereas 60-69 age group has significantly lower risk (19). In addition, there are studies showing that the females are at higher risk of experiencing SAH and its complications (19,21). Our study, however, found no significant gender ($p=0.395$) or age ($p=0.119$) related difference among our patients in terms of risks. Although the studies in the literature generally state that age and gender are important risk factors, there are also articles with findings concurring that of our study (22).

HT is accepted as a risk factor for aneurysm formation and rupture (23). Laboratory studies have shown that a link exists between cerebral aneurysm and HT (24). Our study also found that HT was the most important morbidity accompanying SAH.

The most common site for aneurysm was MCA. In the literature, however, the most common site for aneurysm is ACoA which is the type of aneurysm with the highest rate of mortality and morbidity (3). The mortality rate we found in our study (11.7%) was unaffected by the site of aneurysm. We think that this differentiation may be related to the sample size of our study ($n=44$).

Discussions on the timing of surgery continue and varying views have been put forward about this issue thus far (25-27). However, in our study, we found a statistically significant relationship between the timing of surgery and GOSE score ($p=0.014$): The earlier the surgery was performed, the higher the patient's GOSE values and the lower the mortality and morbidity were.

A strongly negative correlation was found between the HHS and GOSE scores of the patients ($p<0.001$). As HHS score increased, survival rate decreased significantly. We think that it gives a very significant information about the prognosis. Past studies also support our findings (27-30). For example, studies conducted with 100 patients with SAH in 2001, with 3567 patients with SAH in 2007, with 720 patients with SAH in 2020, as well as some other studies conducted with many patients revealed the relationship between HHS score and prognosis (31-33). Like HHS score, WFNS scale score was also found to have a statistically significant relationship with GOSE score ($p=0.002$). As WFNS scale score increased, GOSE score decreased. A look at the literature shows that the findings about WFNS scale score and patient prognosis are compatible with our results (31-36).

In the literature comparing clinical outcome and preoperative parameters, fewer parameters were examined than in our study. We determined that all parameters such as age, gender, day of surgery, HHS, WFNS Scale and Fisher scale scores, and aneurysm localization were not considered.

In our study, it was seen that the findings related to age and gender were not compatible with the literature. Findings on the day of surgery and WFNS scale score were consistent with the literature. Findings on HHS and Fisher scale scores supported the literature.

Study Limitations

The limitations of the study were that it was conducted in a single center, it was retrospective, and the number of patients was low. The low number of patients caused underpowered statistical analysis. Consequently, further comparative, long-term studies with larger patient groups are necessary to confirm these findings.

CONCLUSION

We suggest that WFNS scale and HHS scores of patients with aneurysmal SAH and timing of treatment provide important information about their GOSE scores and thereby the prognosis. Age, gender, comorbidities of the patients and the site of aneurysm did not make a significant contribution to the prognosis of the patients. Identifying and analysing the risk factors in studies with larger sample sizes will contribute to the literature. Conducting the study with a larger patient series will yield more reliable results.

Ethics Committee Approval: Ethical approval was obtained from the University of Health Sciences Turkey, Gaziosmanpaşa Training and Research Hospital Clinical Research Ethics Committee (approval no: 319, date: 22.09.2021).

Informed Consent: Written informed consent for scientific purposes and clinical data collection were obtained from patients according to institutional protocol.

Peer-review: Externally and internally peer-reviewed.

Author Contributions: Surgical and Medical Practices - Ö.Ö.; Concept - Ö.Ö., F.D.; Design - Ö.Ö., O.B.; Data Collection and/or Processing - Ö.Ö., B.B.; Analysis and/or Interpretation - Ö.Ö., S.K.; Literature Search - Ö.Ö.; Writing - Ö.Ö., S.K.

Conflict of Interest: The authors have no conflict of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

Etik Komite Onayı: Sağlık Bilimleri Üniversitesi, Gaziosmanpaşa Eğitim ve Araştırma Hastanesi Klinik Araştırmalar Etik Kurulu'ndan etik onay alındı (onay no: 319, tarih: 22.09.2021).

Hasta Onamı: Bilimsel amaçlar ve klinik veri toplama için yazılı bilgilendirilmiş onam, kurumsal protokole göre hastalardan alınmıştır.

Hakem Değerlendirmesi: Editörler kurulu ve editörler kurulu dışında olan kişiler tarafından değerlendirilmiştir.

Yazar Katkıları: Cerrahi ve Medikal Uygulama - Ö.Ö.; Konsept - Ö.Ö., F.D.; Dizayn - Ö.Ö., O.B.; Veri Toplama veya İşleme - Ö.Ö., B.B.; Analiz veya Yorumlama - Ö.Ö., S.K.; Literatür Arama - Ö.Ö.; Yazan - Ö.Ö., S.K.

Çıkar Çatışması: Yazarlar tarafından çıkar çatışması bildirilmemiştir.

Finansal Destek: Yazarlar tarafından finansal destek almadıkları bildirilmiştir.

REFERENCES

1. Brisman JL, Song JK, Newell DW. Cerebral aneurysms. *N Engl J Med* 2006; 355: 928-39.
2. Ingall TJ, Whisnant JP, Wiebers DO, O'Fallon WM. Has there been a decline in subarachnoid hemorrhage mortality? *Stroke* 1989; 20: 718-24.
3. Brinjikji W, Zhu YQ, Lanzino G, Cloft HJ, Murad MH, Wang Z, et al. Risk Factors for Growth of Intracranial Aneurysms: A Systematic Review and Meta-Analysis. *AJNR Am J Neuroradiol* 2016; 37: 615-20.
4. Rincon F, Rossenwasser RH, Dumont A. The epidemiology of admissions of nontraumatic subarachnoid hemorrhage in the United States. *Neurosurgery* 2013; 73: 217-22; discussion 212-3.
5. Bradac O, Hide S, Mendelow DA, Benes V. Aneurysm treatment in Europe 2010: an internet survey. *Acta Neurochir (Wien)* 2012; 154: 971-8; discussion 977-8.
6. Wiebers DO, Whisnant JP, Huston J 3rd, Meissner I, Brown RD Jr, Piepgras DG, et al. Unruptured intracranial aneurysms: natural history, clinical outcome, and risks of surgical and endovascular treatment. *Lancet* 2003; 362: 103-10.
7. Molyneux A, Kerr R, Stratton I, Sandercock P, Clarke M, Shrimpton J, et al. International Subarachnoid Aneurysm Trial (ISAT) of neurosurgical clipping versus endovascular coiling in 2143 patients with ruptured intracranial aneurysms: a randomised trial. *Lancet* 2002; 360: 1267-74.
8. Taylor CJ, Robertson F, Brealey D, O'shea F, Stephen T, Brew S, et al. Outcome in poor grade subarachnoid hemorrhage patients treated with acute endovascular coiling of aneurysms and aggressive intensive care. *Neurocrit Care* 2011; 14: 341-7.
9. Seule M, Oswald D, Muroi C, Brandi G, Keller E. Outcome, Return to Work and Health-Related Costs After Aneurysmal Subarachnoid Hemorrhage. *Neurocrit Care* 2020; 33: 49-57.
10. Schwartz C, Pfeifferkorn T, Ebrahimi C, Ottomeyer C, Fesl G, Bender A, et al. Long-term Neurological Outcome and Quality of Life after World Federation of Neurosurgical Societies Grades IV and V Aneurysmal Subarachnoid Hemorrhage in an Interdisciplinary Treatment Concept. *Neurosurgery* 2017; 80: 967-74.
11. Rosen DS, Macdonald RL. Subarachnoid hemorrhage grading scales: a systematic review. *Neurocrit Care* 2005; 2: 110-8.
12. Oshiro EM, Walter KA, Piantadosi S, Witham TF, Tamargo RJ. A new subarachnoid hemorrhage grading system based on the Glasgow Coma Scale: a comparison with the Hunt and Hess and World Federation of Neurological Surgeons Scales in a clinical series. *Neurosurgery* 1997; 41: 140-7; discussion 147-8.
13. Wilson JT, Pettigrew LE, Teasdale GM. Structured interviews for the Glasgow Outcome Scale and the extended Glasgow Outcome Scale: guidelines for their use. *J Neurotrauma* 1998; 15: 573-85.
14. de Rooij NK, Linn FH, van der Plas JA, Algra A, Rinkel GJ. Incidence of subarachnoid haemorrhage: a systematic review with emphasis on region, age, gender and time trends. *J Neurol Neurosurg Psychiatry* 2007; 78: 1365-72.
15. Roganović Z, Pavličević G. Factors influencing the outcome after the operative treatment of cerebral aneurysms of anterior circulation. *Vojnosanit Pregl* 2002; 59: 463-71.
16. Cánovas D, Gil A, Jato M, de Miquel M, Rubio F. Clinical outcome of spontaneous non-aneurysmal subarachnoid hemorrhage in 108 patients. *Eur J Neurol* 2012; 19: 457-61.
17. Fontanella M, Rainero I, Panciani PP, Schatlo B, Benevello C, Garbossa D, et al. follow-up. *Neurosurg Rev* 2011; 34: 477-84.
18. Ellis JA, McDowell MM, Mayer SA, Lavine SD, Meyers PM, Connolly ES Jr. The role of antiplatelet medications in angiogram-negative subarachnoid hemorrhage. *Neurosurgery* 2014; 75: 530-5; discussion 534-5.
19. Malinova V, Schatlo B, Voit M, Suntheim P, Rohde V, Mielke D. Identification of specific age groups with a high risk for developing cerebral vasospasm after aneurysmal subarachnoid hemorrhage. *Neurosurg Rev* 2016; 39: 429-36.
20. Kale SP, Edgell RC, Alsheklee A, Borhani Haghighi A, Sweeny J, Felton J, et al. Age-associated vasospasm in aneurysmal subarachnoid hemorrhage. *J Stroke Cerebrovasc Dis* 2013; 22: 22-7.

21. Knekt P, Reunanen A, Aho K, Heliövaara M, Rissanen A, Aromaa A, et al. Risk factors for subarachnoid hemorrhage in a longitudinal population study. *J Clin Epidemiol* 1991; 44: 933-9.
22. Lindbohm JV, Kaprio J, Jousilahti P, Salomaa V, Korja M. Sex, Smoking, and Risk for Subarachnoid Hemorrhage. *Stroke* 2016; 47: 1975-81.
23. Connolly ES Jr, Rabinstein AA, Carhuapoma JR, Derdeyn CP, Dion J, Higashida RT, et al. Guidelines for the management of aneurysmal subarachnoid hemorrhage: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2012; 43: 1711-37.
24. Morimoto M, Miyamoto S, Mizoguchi A, Kume N, Kita T, Hashimoto N. Mouse model of cerebral aneurysm: experimental induction by renal hypertension and local hemodynamic changes. *Stroke* 2002; 33: 1911-5.
25. Kassell NF, Torner JC, Haley EC Jr, Jane JA, Adams HP, Kongable GL. The International Cooperative Study on the Timing of Aneurysm Surgery. Part 1: Overall management results. *J Neurosurg* 1990; 73: 18-36.
26. Torner JC, Kassell NF, Wallace RB, Adams HP Jr. Preoperative prognostic factors for rebleeding and survival in aneurysm patients receiving antifibrinolytic therapy: report of the Cooperative Aneurysm Study. *Neurosurgery* 1981; 9: 506-13.
27. Niskanen MM, Hernesniemi JA, Vapalahti MP, Kari A. One-year outcome in early aneurysm surgery: prediction of outcome. *Acta Neurochir (Wien)* 1993; 123: 25-32.
28. Bansal M, Mittal RS. Clinical Demographic Association and Outcome in Patients with Aneurysmal Subarachnoid Hemorrhage. *Indian Journal of Neurosurgery* 2015; 4: 63-8.
29. Bhat AR, Afzalwani M, Kirmani AR. Subarachnoid hemorrhage in Kashmir: Causes, risk factors, and outcome. *Asian J Neurosurg* 2011; 6: 57-71.
30. Sodhi HB, Savardekar AR, Mohindra S, Chhabra R, Gupta V, Gupta SK. The clinical profile, management, and overall outcome of aneurysmal subarachnoid hemorrhage at the neurosurgical unit of a tertiary care center in India. *J Neurosci Rural Pract* 2014; 5: 118-26.
31. Bonilha L, Marques EL, Carelli EF, Fernandes YB, Cardoso AC, Maldaum MV, et al. Risk factors and outcome in 100 patients with aneurysmal subarachnoid hemorrhage. *Arq Neuropsiquiatr* 2001; 59: 676-80.
32. Rosengart AJ, Schultheiss KE, Tolentino J, Macdonald RL. Prognostic factors for outcome in patients with aneurysmal subarachnoid hemorrhage. *Stroke* 2007; 38: 2315-21.
33. Chen YB, Wang Z, Zhou D, Zhang SM, Yu ZQ, Chen G. [Analysis of prognosis and risk factors of 730 patients with anterior communicating aneurysm]. *Zhonghua Yi Xue Za Zhi* 2020; 100: 3515-9. Chinese.
34. Aydın Y, Halit Çavuşoğlu H. Subarachnoid hemorrhage. *Med Bull Sisli Etfal Hosp* 2007; 41: 7-16.
35. Uysal E, Kırdar Ö, Özel A, Ertürk ŞM, Yıldırım H, Caymaz I, et al. Ecstasy use complicating with ICA dissection; A case report. *Med Bull Sisli Etfal Hosp* 2008; 42: 23-5.
36. Uysal E, Ulusay SM, Civelek Ş, Başak M. Combined endovascular and surgery treatment for neck and head paragangliomas. *Med Bull Sisli Etfal Hosp* 2010; 44: 27-31.