

The Effects of the COVID-19 Pandemic on Patients with Trauma Presented to the Emergency Department: A Multicentre Experience in İstanbul

COVID-19 Pandemisinin Acil Servise Başvuran Travmalı Hastalar Üzerindeki Etkileri: İstanbul'da Çok Merkezli Bir Deneyim

Mustafa Çalık¹, Dilay Satılmış², Burcu Genç Yavuz³, Ramazan Güven⁴, Şahin Çolak³, Özgür Söğüt⁵, İbrahim Altundağ³, Ertuğrul Altınbilek⁶, İsmail Tayfur⁷, Nihat Müjdat Hökenek⁸, Gökhan Eyüpoğlu⁹, Tarık Akdemir⁵

¹University of Health Sciences Turkey, Gaziosmanpaşa Training and Research Hospital, Clinic of Emergency Medicine, İstanbul, Turkey

²University of Health Sciences Turkey, Sultan 2. Abdülhamid Han Training and Research Hospital, Clinic of Emergency Medicine, İstanbul, Turkey

³University of Health Sciences Turkey, Haydarpaşa Numune Training and Research Hospital, Clinic of Emergency Medicine, İstanbul, Turkey

⁴University of Health Sciences Turkey, Kanuni Sultan Süleyman Training and Research Hospital, Clinic of Emergency Medicine, İstanbul, Turkey

⁵University of Health Sciences Turkey, İstanbul Haseki Training and Research Hospital, Clinic of Emergency Medicine, İstanbul, Turkey

⁶University of Health Sciences Turkey, Şişli Hamidiye Etfal Training and Research Hospital, Clinic of Emergency Medicine, İstanbul, Turkey

⁷University of Health Sciences Turkey, Sancaktepe Şehit Prof. Dr. İlhan Varank Training and Research Hospital, Clinic of Emergency Medicine, İstanbul, Turkey

⁸University of Health Sciences Turkey, Kartal Dr. Lütfi Kırdar City Hospital, Clinic of Emergency Medicine, İstanbul, Turkey

⁹University of Health Sciences Turkey, Başakşehir Çam and Sakura City Hospital, Clinic of Emergency Medicine, İstanbul, Turkey

Cite this article as: Çalık M, Satılmış D, Genç Yavuz B, Güven R, Çolak Ş, Söğüt Ö, Altundağ İ, Altınbilek E, Tayfur İ, Hökenek NM, Eyüpoğlu G, Akdemir T. The Effects of the COVID-19 Pandemic on Patients with Trauma Presented to the Emergency Department: A Multicentre Experience in İstanbul. J Acad Res Med 2023;13(1):30-5

ABSTRACT

Objective: This study examines the characteristics of patients with trauma who presented to the emergency department before and after the novel coronavirus disease-2019 (COVID-19) pandemic and analyze the effects on the traumatic injuries.

Methods: In this study, medical records of patients who presented to the emergency departments of seven hospitals operating as tertiary level hospitals in İstanbul between March-June 2018, March-June 2019, and March-June 2020 are due to trauma were retrospectively analyzed. The clinical and demographic characteristics of traumatic injuries before and after the pandemic were compared.

Results: In our study, 4088 trauma patients' data were reviewed: 1279 in March-June 2018, 1684 in March-June 2019, and 1125 in March-June 2020. When the total number of patients was examined, it was noticed that the number of patients decreased significantly during the COVID-19 pandemic. No significant difference was found between the periods regarding the sex and trauma mechanisms of the patients. The mean age was higher in patients admitted in 2020 compared to previous years. In our study, incidences of intracranial hemorrhage, femur fracture, lung injuries, and mortality rates were higher in March-June 2020 compared to previous years.

Conclusion: In March-June 2020, compared to the previous year, there was a 34% decrease in trauma cases admitted to the emergency department. Albeit no difference was found between the periods regarding trauma mechanisms, the higher mortality in the March-June 2020 period indicates that trauma continues to be one of the major causes of death despite the pandemic.

Keywords: COVID-19, emergency, injuries, trauma

ORCID IDs of the authors: M.Ç. 0000-0002-3184-2943; D.S. 0000-0003-3765-2208; B.G.Y. 0000-0001-6693-5288; R.G. 0000-0003-4129-8985; Ş.Ç. 0000-0001-8192-9652; Ö.S. 0000-0003-3365-3713; İ.A. 0000-0002-0880-7218; E.A. 0000-0003-4201-8850; İ.T. 0000-0002-4852-7915; N.M.H. 0000-0002-8780-572X; G.E. 0000-0001-8583-3690; T.A. 0000-0003-1178-2274.

Corresponding Author/Sorumlu Yazar: Mustafa Çalık,

E-mail: drmustafacalik@yahoo.com

Received Date/Geliş Tarihi: 24.03.2023 **Accepted Date/Kabul Tarihi:** 31.03.2023

©Copyright 2023 by University of Health Sciences Turkey Gaziosmanpaşa Training and Research Hospital.

Journal of Academic Research in Medicine published by Galenos Publishing House.

Licensed by Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International

(CC BY-NC-ND 4.0) Available on-line at www.jarem.org



ÖZ

Amaç: Bu çalışmanın amacı, yeni koronavirus hastalığı-2019 (COVID-19) salgını öncesi ve sonrası paralel dönemlerde acil servise başvuran travmalı hastaların özelliklerini incelemek ve toplumsal değişimin travmatik yaralanmalar üzerindeki etkilerini analiz etmektir.

Yöntemler: Bu çalışmada, Mart-Haziran 2018, Mart-Haziran 2019 ve Mart-Haziran 2020 tarihleri arasında İstanbul'da üçüncü basamak eğitim ve araştırma hastanesi olarak faaliyet gösteren yedi hastanenin acil servislerine travma nedeniyle başvuran hastaların tıbbi kayıtları retrospektif olarak incelenmiştir. Pandemi öncesi ve sonrası travmatik yaralanmaların klinik ve demografik özellikleri karşılaştırıldı.

Bulgular: Çalışmamızda 4088 travma hastasının verileri incelendi; Mart-Haziran 2018'de 1279, Mart-Haziran 2019'da 1684 ve Mart-Haziran 2020'de 1125. Toplam hasta sayısı incelendiğinde, COVID-19 pandemisi sırasında hasta sayısının önemli ölçüde azaldığı fark edildi. Hastaların cinsiyet ve travma mekanizmaları açısından dönemler arasında anlamlı fark bulunmadı. 2020 yılında başvuran hastalarda yaş ortalaması önceki yıllara göre daha yüksekti. Çalışmamızda Mart-Haziran 2020 döneminde kafa içi kanama, femur kırığı, akciğer yaralanması insidansı ve mortalite oranları önceki paralel yıllara göre daha yüksekti.

Sonuç: Mart-Haziran 2020 döneminde bir önceki yıla göre acil servise başvuran travma olgularında %34 azalma oldu. Travma mekanizmaları açısından dönemler arasında fark bulunmasa da Mart-Haziran 2020 döneminde ölüm oranlarının yüksek olması, pandemiye rağmen travmanın en önemli ölüm nedenlerinden biri olmaya devam ettiğini gösteriyor.

Anahtar kelimeler: COVID-19, acil, yaralanmalar, travma

INTRODUCTION

Despite increased trauma prevention measures and advances in post-trauma care, trauma-related deaths remain one of the main causes of mortality worldwide (1). Likewise, it has been reported that roughly 3% of all deaths in our country are due to trauma, and the most common cause of these deaths is motor vehicle accidents (2).

The coronavirus disease-2019 (COVID-19) pandemic, which broke out in December 2019 and spread worldwide in a short time, still continues to affect millions of people. Morbidity and mortality are attempted to be prevented through treatment strategies and widespread vaccination. Upon the declaration of COVID-19 as a pandemic by the World Health Organization, measures, such as lockdown, social isolation, or restriction of crowded organizations, were taken in many countries. Although the measures were reduced from time to time, these measures were repeated at the peak times of the outbreak. Social restrictions also decreased hospital admissions due to non-COVID-19 causes (3,4). In the literature, it has been reported that there is a decrease in acute coronary syndrome (5), stroke (6), orthopedic (7,8), and pediatric trauma (9) cases during the pandemic.

Similarly, in our country, it was seen that outpatient clinic applications other than COVID-19 decreased, and elective surgeries were postponed due to social isolation measures (3,10). However, the patient profile of the emergency departments diversified with the addition of COVID-19 cases. Although studies for certain disease prevalence have been revealed in the literature in this process, our study compares the demographic characteristics of patients with trauma who presented to seven different hospitals before and after the COVID-19 pandemic.

METHODS

This was a retrospective, multicenter, cross-sectional study, including 4088 patients with trauma who were admitted to University of Health Sciences Turkey, İstanbul Gaziosmanpaşa

Training and Research Hospital, Haydarpaşa Numune Training and Research Hospital, Kanuni Sultan Süleyman Training and Research Hospital, İstanbul Haseki Training and Research Hospital, Şişli Hamidiye Etfal Training and Research Hospital, Sancaktepe Şehit Prof. Dr. İlhan Varank Training and Research Hospital, and Kartal Dr. Lütfi Kırdar City Hospital Emergency Department. These seven centers where data was collected were tertiary trauma centers in İstanbul, the most populous city of Turkey, to which all age groups applied. Before conducting this study, ethical approval was obtained from the University of Health Sciences Turkey, Kanuni Sultan Süleyman Training and Research Hospital Clinical Research Ethics Committee (decision no: 133, date: 08.07.2020).

In our study, the period before and during the pandemic was divided into three periods as March-June 2018, March-June 2019, and March-June 2020. March-June 2018 and 2019 were the period when COVID-19 cases were not yet identified worldwide, while March-June 2020 was the period when a pandemic was declared worldwide and our country went into total closure. The medical records of the patients admitted to the above-mentioned centers due to trauma on these dates were reviewed retrospectively. Our inclusion criterion was to be applied to the emergency department after having acute trauma in any age group. If the same patient presented to the emergency department more than once in the specified date range, only the first admission was taken into consideration. Patients who were transferred from another hospital to surgery, orthopedics, or another branch due to trauma were excluded from this study.

Age, sex, trauma mechanism, discharge, or hospitalization, and traumatic injuries of the patients were noted down from the hospital records, and the changes in trauma admissions during the pandemic were analyzed.

The primary outcome of our study was to analyze the effects of the COVID-19 pandemic on patients with trauma presented to the emergency department and its effects on trauma-related injuries. Our secondary outcome was to investigate the effects of the pandemic on traumatic deaths and discharges.

Statistical Analysis

Continuous data were presented with mean \pm standard deviation and with frequency (n) and percentage (%) for the categorical data. The normality assumptions were controlled by the Shapiro-Wilk test. The association between categorical data was determined by Pearson chi-square and Fisher's Exact test. One-Way ANOVA was used to compare age among study groups, and the Tukey's HSD (honestly significant difference) test was used as a post-hoc test for significant cases. Statistical analysis was conducted using the software of IBM SPSS Statistics for Windows, Version 23.0 (IBM Corp., Armonk, NY). A two-sided p-value less than 0.05 was considered statistically significant.

RESULTS

In our study, the data of 4088 patients with trauma, 2963 in the pre-COVID-19 period and 1125 in the March-June 2020 period, were analyzed. The mean age of patients who presented in March-June 2018 was 42.6 ± 25.9 years, while it was 41.4 ± 25.7 years for those in March-June 2019, and 44.1 ± 25.1 years for those who applied in March-June 2020. The mean age of the patients who presented in 2020 was significantly higher than in 2019 ($p=0.021$) (Table 1). When we examined the number of trauma patients in the March-June periods of 2018, 2019, and 2020, it was noticed

that the number of cases during the pandemic decreased by 34% compared to 2019. No significant difference was determined regarding the sex distribution of the patients by years ($p=0.282$). When we examined the trauma mechanisms according to 2018, 2019, and 2020, no significant difference was found between the trauma mechanisms of the patients who were admitted to the hospital in the same months before and during the pandemic (Table 2).

The incidence of finger amputation in 2020 (1.2%) was higher than in 2019 (0.4%) ($p=0.034$), whereas the incidence of shoulder dislocation in 2020 (0.5%) was higher than in 2018 (0%) ($p=0.022$), and the incidence of other joint dislocations in 2018 (0.6%) was significantly higher than in 2020 (0%) ($p=0.014$). Motor accident ($p=0.056$), occupational accidents ($p=0.095$), and foot sprain ($p=0.073$) incidences were higher in 2020, and the incidences of falling off a bicycle ($p=0.072$) in 2019 were higher, yet the difference was not significant.

In our study, the incidence of intracranial hemorrhage in 2020 (13%) was higher than in 2018 (9.7%) ($p=0.021$), whereas the incidence of lumbar vertebral fracture in 2019 (2.8%) was higher than in 2020 (1.4%) ($p=0.049$). The incidence of hemothorax in 2020 (3.1%) was higher than that in 2018 (1.6%) and 2019 (1.8%) ($p=0.026$). The lung contusion incidence in 2020 was 5.6%, which was higher than in

Table 1. Demographic characteristics

Demographics	2018 (n=1279)	2019 (n=1684)	2020 (n=1125)	p-value
Sex, n (%)				
Female	470 (36.7)	581 (34.5)	381 (33.9)	0.282
Male	809 (63.3)	1103 (65.5)	744 (66.1)	
Age (years), mean \pm SD	$42.6 \pm 25.9^{a,b}$	41.4 ± 25.7^a	44.1 ± 25.1^b	0.021

SD: standard deviation, HSD: honestly significant difference

Pearson chi-square test. One-Way ANOVA with post-hoc Tukey HSD test. Same letters in a row denote the lack of statistically significant difference

Table 2. Mechanism of injury

Mechanism, n (%)	2018 (n=1279)	2019 (n=1684)	2020 (n=1125)	p-value
Motor vehicle accident	102 (8)	110 (6.5)	74 (6.6)	0.254
Auto versus pedestrian	190 (14.9)	243 (14.4)	149 (13.2)	0.507
Motorcycle accident	61 (4.8)	106 (6.3)	79 (7)	0.056
Fall from high place	125 (9.8)	178 (10.6)	113 (10)	0.766
Stab wound	48 (3.8)	62 (3.7)	50 (4.4)	0.557
Firearm injuries	32 (2.5)	58 (3.4)	42 (3.7)	0.189
Bicycle accident	23 (1.8)	33 (2)	10 (0.9)	0.072
Burn	8 (0.6)	4 (0.2)	8 (0.7)	0.127
Blunt assault	56 (4.4)	61 (3.6)	45 (4)	0.578
Industrial injury	34 (2.7)	65 (3.9)	47 (4.2)	0.095
Ingress of foreign body	12 (0.9)	17 (1)	14 (1.2)	0.745
Limb impingement	1 (0.1)	2 (0.1)	2 (0.2)	0.863
Suicide	2 (0.2)	0 (0)	0 (0)	0.174

Pearson chi-square test, Fisher's Exact test

2018 (1.3%) and 2019 (2.6%) ($p<0.001$). The incidence of intestinal perforation (1.3%) was significantly higher in 2018 than 2019 (0.1%) and 2020 (0.4%) ($p<0.001$). While the incidence of urinary bladder injury in 2020 (0.7%) was higher than in 2018 (0.1%) ($p=0.033$), the incidence of intra-abdominal fluid in 2020 (4.1%) was higher than in 2018 (2.1%) and 2019 (2%) ($p=0.001$), the soft tissue injury rate in 2018 (15.8%) was higher ($p=0.021$) than in 2019 (12.4%), and the incidence of simple cuts or abrasions (17.9%) in 2020 was higher than in 2018 (12.2%) and 2019 (12.5%) ($p<0.001$). When the distribution of fracture types by years was analyzed, the incidence of costal

fracture (3.6%) in 2019 and 2020 was higher ($p<0.001$) than in 2018 ($p<0.001$), whereas the rate of radius fracture and hand phalanx fracture (8.6% and 3.6%) in 2019 were higher ($p=0.005$ and $p=0.030$, respectively) than in 2020 (5.5% and 2%). The incidence of femoral fractures in 2020 (22%) was higher than in 2019 (17.5%) ($p=0.009$). The incidence of muscle tears in 2019 (3.1%) was significantly higher than in 2018 (1.6%), whereas the rate of other tears in 2018 (9.4%) was significantly higher than in 2019 (5.5%) and 2020 (6%) ($p=0.011$ and $p<0.001$) (Table 3). The rate of patients who died in 2020 (4.3%) was significantly higher than in 2018 (2.1%) and 2019 (2.4%)

Table 3. Diagnoses per year

Main diagnosis, n (%)	2018 (n=1279)	2019 (n=1684)	2020 (n=1125)	p-value
Cranial fracture	138 (10.8)	204 (12.1)	116 (10.3)	0.283
Intracranial hemorrhage	124 (9.7) ^a	172 (10.2) ^{a,b}	146 (13) ^b	0.021
Cervical vertebral fracture	11 (0.9)	9 (0.5)	13 (1.2)	0.191
Thoracic vertebral fracture	24 (1.9)	29 (1.7)	18 (1.6)	0.873
Lumbar vertebral fracture	27 (2.1) ^{a,b}	47 (2.8) ^a	16 (1.4) ^b	0.049
Thoracic injury	39 (3) ^a	82 (4.9) ^b	87 (7.7) ^c	<0.001
Hemothorax	21 (1.6) ^a	31 (1.8) ^a	35 (3.1) ^b	0.026
Pneumothorax	25 (2)	54 (3.2)	31 (2.8)	0.112
Lung contusion	17 (1.3) ^a	43 (2.6) ^a	63 (5.6) ^b	<0.001
Costal fracture	13 (1) ^a	60 (3.6) ^b	41 (3.6) ^b	<0.001
Cardiac injury	5 (0.4)	2 (0.1)	0 (0)	0.064
Great vessel injury	8 (0.6)	11 (0.7)	6 (0.5)	0.936
Internal organ injury	56 (4.4)	61 (3.6)	52 (4.6)	0.371
Liver injury	18 (1.4)	16 (1)	10 (0.9)	0.379
Spleen injury	11 (0.9)	19 (1.1)	9 (0.8)	0.624
Intestinal perforation	16 (1.3) ^a	1 (0.1) ^b	4 (0.4) ^b	<0.001
Kidney injury	8 (0.6)	7 (0.4)	2 (0.2)	0.236
Urinary bladder injury	1 (0.1) ^a	5 (0.3) ^{a,b}	8 (0.7) ^b	0.033
Intra-abdominal free fluid	27 (2.1) ^a	34 (2) ^a	46 (4.1) ^b	0.001
Simple incisions/graze	156 (12.2) ^a	211 (12.5) ^a	201 (17.9) ^b	<0.001
Orthopedic fracture	910 (71.1)	1231 (73.1)	813 (72.3)	0.502
Humeral fracture	103 (8.1)	162 (9.6)	97 (8.6)	0.314
Ulna fracture	70 (5.5)	76 (4.5)	44 (3.9)	0.182
Radius fracture	83 (6.5) ^{a,b}	144 (8.6) ^a	62 (5.5) ^b	0.005
Metacarpal bone fracture	28 (2.2)	33 (2)	29 (2.6)	0.549
Hand phalanx fracture	30 (2.3) ^{a,b}	60 (3.6) ^a	23 (2) ^b	0.030
Pelvic fracture	98 (7.7)	134 (8)	86 (7.6)	0.938
Femur fracture	258 (20.2) ^{a,b}	294 (17.5) ^a	248 (22) ^b	0.009
Tibia fracture	182 (14.2)	252 (15)	162 (14.4)	0.837
Fibula fracture	71 (5.6)	112 (6.7)	87 (7.7)	0.099
Metatarsal fracture	42 (3.3)	49 (2.9)	45 (4)	0.286
Foot phalanx fracture	20 (1.6)	27 (1.6)	21 (1.9)	0.819
Muscle tears	20 (1.6) ^a	53 (3.1) ^b	22 (2) ^{a,b}	0.011
Other fractures	120 (9.4) ^a	92 (5.5) ^b	68 (6) ^b	<0.001

Pearson chi-square test, Fisher's Exact test. Same letters in a row denote the lack of statistically significant difference

($p=0.003$). In 4052 patients who were negative for COVID-19, the exitus rate (3.6%) in 2020 was higher than in 2018 (2.1%) ($p=0.037$). While 72.2% of the 36 patients with COVID-19 polymerase chain reaction (PCR) positive were discharged and 2.8% were referred, 25% were exitus (Table 4).

DISCUSSION

In our study, we determined that there was a 34% decrease in trauma cases admitted to the emergency department in the March-June 2020 period compared to the previous year. Besides, no significant difference was determined between the mentioned periods regarding trauma mechanisms. However, we found a higher post-traumatic death rate during the pandemic.

The decrease in trauma cases during the pandemic period may be due to the decrease in traumatic injuries because of the social restrictions and lockdown mandates, or because patients after minor trauma did not apply to hospitals for fear of COVID-19 contamination. Similar to our study, Hartnett et al. (11) compared March 29-April 25, 2020 with March 31-April 27, 2019, and found that there was a 42% decrease in the department of emergency admissions. İlhan et al. (12), in their single-center study, revealed a 60% decrease in trauma admissions during the pandemic. In another study compared before and after lockdown, it was shown that post-traumatic hospitalization decreased by 30.9% during the lockdown period (13). In another study conducted with 618 patients, it was stated that the need for open surgery increased in the pandemic period compared with the pre-pandemic period, although a 25% decrease was detected in the patients who applied to the emergency department for surgical reasons during the pandemic period (10). In the study investigating the injury-related hospitalization of 21 hospitals between March 15-April 30, 2016-2020, it was shown that hospitalization decreased by 26% during the lockdown period (14).

Regarding the trauma mechanisms, Chiba et al. (15) found a 38.7% reduction in motorcycle accidents, a 42.5% reduction in auto versus pedestrian accidents, and a 28.4% reduction in bicycle accidents. However, they did not report a significant reduction in motor vehicle accidents during the pandemic (15). Besides, DiFazio et al. (16) reported a 75% reduction in motor vehicle accidents and a 28.9% reduction in falls, along with a 44.9% reduction in trauma admissions during the lockdown period of the pandemic. Moreover, İlhan et al. (12) detected a significant decrease in pedestrian injuries during the pandemic. On the other hand, in our study, unlike these studies, the trauma mechanism was similar in the specified periods of all three years. Albeit a significant decrease in traumatic injuries was an expected finding during the pandemic, it was remarkable for us that the trauma mechanism did not change. The reason for this may be that although there are social restrictions and lockdown, a substantial group, such as people with special work permits or motor couriers, is exempt from these bans. Additionally, although the decrease in the number of vehicles on the road relieves Istanbul traffic, the trauma mechanism may not change compared to previous years due to the use of vehicles at higher speeds on relatively empty roads. When we examined the traumatic injuries in our study, the fact that intracranial bleeding, hemothorax, lung contusion, intestinal perforation, urinary system injury, free fluid in the abdomen, soft tissue injury, rib fracture, and femur fracture were detected more frequently in the March-June 2020 period compared to previous years support the exposure to high-energy trauma during the pandemic period. Hence, we consider that the enforcement of traffic rules and raising public awareness about traumas are of considerable importance even during lockdown periods.

Chiba et al. (15) revealed that mortality decreased during the lockdown period compared with the previous year. On the other

Table 4. Recent outcome of patients by year

Outcome n (%)	2018 (n=1279)	2019 (n=1684)	2020 (n=1125)	p-value
All patients (n=4088)				
Discharge	1238 (96.8) ^a	1633 (97) ^a	1072 (95.3) ^a	0.003
Referral to another hospital	14 (1.1) ^a	10 (0.6) ^a	5 (0.4) ^a	
Exitus	27 (2.1) ^a	41 (2.4) ^a	48 (4.3) ^b	
COVID-19 PCR negative patients (n=4052)				
Discharge	1238 (96.8) ^a	1633 (97) ^a	1046 (96.1) ^a	0.037
Referral to another hospital	14 (1.1) ^a	10 (0.6) ^a	4 (0.4) ^a	
Exitus	27 (2.1) ^a	41 (2.4) ^{a,b}	39 (3.6) ^b	
COVID-19 PCR positive patients (n=36)				
Discharge	-	-	26 (72.2)	-
Referral to another hospital	-	-	1 (2.8)	
Exitus	-	-	9 (25)	

COVID-19: coronavirus disease-2019, PCR: polymerase chain reaction
Pearson chi-square test, same letters in a row denote the lack of statistically significant difference

hand, Yasin et al. (17) showed that mortality was higher during the pandemic period in their study, in which they included 750 road traffic collision patients during the pre-COVID-19 period and 499 road traffic collision patients during the COVID-19 period. Likewise, when we looked at the post-traumatic death rates in our study, there was a significant increase in the death rate in 2020. The reason for this may be the occurrence and mortality of high-energy traumatic injuries at higher speeds in reduced traffic. Furthermore, a more elective approach to the patients due to the pandemic and the waiting for the PCR results for the operation during the pandemic might increase the mortality.

Study Limitations

The current study has some limitations. First, it was a retrospective study and therefore, some data were not available. Trauma scores could not be calculated due to missing data. Although we showed that some injuries were more common in the COVID-19 period in our study, we could not compare the trauma scores with the pre- and post-COVID-19 periods. Second, although we conducted a multicenter study, regional studies with larger participation are needed to analyze the effects of the COVID-19 pandemic and social isolation on trauma.

CONCLUSION

In line with the literature, lockdown and social isolation measures mandated in many countries due to the pandemic resulted in a remarkable decrease in the number of traumas admitted to hospitals in our country. However, when we examined the trauma mechanism during the lockdown, there was no significant change compared with the previous years, but the mortality rate was higher, especially with intracranial hemorrhage, femur fracture, and lung injuries. These findings indicate that trauma remains one of the major causes of death despite the pandemic. Thus, it supports the significance of planning trauma care and necessary measurements in today's peak day of COVID-19 and a likely new lockdown period to prevent mortality.

Ethics Committee Approval: Before conducting this study, ethical approval was obtained from the University of Health Sciences Turkey, Kanuni Sultan Süleyman Training and Research Hospital Clinical Research Ethics Committee (decision no: 133, date: 08.07.2020).

Informed Consent: Retrospective study.

Peer-review: Externally and internally peer-reviewed.

Author Contributions: Concept - M.Ç., B.G.Y., R.G.; Design - M.Ç., R.G.; Data Collection and/or Processing - M.Ç., D.S., B.G.Y., R.G., Ş.Ç., Ö.S., İ.A., E.A., İ.T., N.M.H., G.E., T.A.; Analysis and/or Interpretation - D.S., B.G.Y., R.G., İ.T.; Literature Search - M.Ç., B.G.Y.; Writing - B.G.Y., Ş.Ç., E.A.

Conflict of Interest: The authors have no conflict of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

Etik Komite Onayı: Bu çalışma yapılmadan önce Sağlık Bilimleri Üniversitesi, Kanuni Sultan Süleyman Eğitim ve Araştırma Hastanesi Klinik Araştırmalar Etik Kurulu'ndan etik onay alınmıştır (karar no: 133, tarih: 08.07.2020).

Hasta Onamı: Retrospektif çalışma.

Hakem Değerlendirmesi: Editörler kurulu ve editörler kurulu dışında olan kişiler tarafından değerlendirilmiştir.

Yazar Katkıları: Konsept - M.Ç., B.G.Y., R.G.; Dizayn - M.Ç., R.G.; Veri Toplama veya İşleme - M.Ç., D.S., B.G.Y., R.G., Ş.Ç., Ö.S., İ.A., E.A., İ.T., N.M.H., G.E., T.A.; Analiz veya Yorumlama - D.S., B.G.Y., R.G., İ.T.; Literatür Arama - M.Ç., B.G.Y.; Yazan - B.G.Y., Ş.Ç., E.A.

Çıkar Çatışması: Yazarlar tarafından çıkar çatışması bildirilmemiştir.

Finansal Destek: Yazarlar tarafından finansal destek almadıkları bildirilmiştir.

REFERENCES

1. Kamine TH, Rembisz A, Barron RJ, Baldwin C, Kromer M. Decrease in Trauma Admissions with COVID-19 Pandemic. *West J Emerg Med* 2020; 21: 819-22.
2. Orhon R, Eren SH, Karadayı S, Korkmaz I, Coşkun A, Eren M, et al. Comparison of trauma scores for predicting mortality and morbidity on trauma patients. *Ulus Travma Acil Cerrahi Derg* 2014; 20: 258-64.
3. İlhan B, Bozdereli Berikol G, Dogan H. Impact of COVID-19 Outbreak on Emergency Visits and Emergency Consultations: A Cross-Sectional Study. *Cureus* 2021; 13: e14052.
4. Cano-Valderrama O, Morales X, Ferrigni CJ, Martín-Antona E, Turrado V, García A, et al. Acute Care Surgery during the COVID-19 pandemic in Spain: Changes in volume, causes and complications. A multicentre retrospective cohort study. *Int J Surg* 2020; 80: 157-61.
5. Mafham MM, Spata E, Goldacre R, Gair D, Curnow P, Bray M, et al. COVID-19 pandemic and admission rates for and management of acute coronary syndromes in England. *Lancet* 2020; 396: 381-9.
6. Bres Bullrich M, Fridman S, Mandzia JL, Mai LM, Khaw A, Vargas Gonzalez JC, et al. COVID-19: Stroke Admissions, Emergency Department Visits, and Prevention Clinic Referrals. *Can J Neurol Sci* 2020; 47: 693-6.
7. Greenhalgh M, Dupley L, Unsworth R, Boden R. Where did all the trauma go? A rapid review of the demands on orthopaedic services at a UK Major Trauma Centre during the COVID-19 pandemic. *Int J Clin Pract* 2021; 75: e13690.
8. Gumina S, Proietti R, Polizzotti G, Carbone S, Candela V. The impact of COVID-19 on shoulder and elbow trauma: an Italian survey. *J Shoulder Elbow Surg* 2020; 29: 1737-42.
9. Nabian MH, Vosoughi F, Najafi F, Khabiri SS, Nafisi M, Veisi J, et al. Epidemiological pattern of pediatric trauma in COVID-19 outbreak: Data from a tertiary trauma center in Iran. *Injury* 2020; 51: 2811-5.
10. Göksoy B, Akça MT, Inanç ÖF. The impacts of the COVID-19 outbreak on emergency department visits of surgical patients. *Ulus Travma Acil Cerrahi Derg* 2020; 26: 685-92.
11. Hartnett KP, Kite-Powell A, DeVies J, Coletta MA, Boehmer TK, Adjemian J, et al. Impact of the COVID-19 Pandemic on Emergency Department Visits - United States, January 1, 2019-May 30, 2020. *MMWR Morb Mortal Wkly Rep* 2020; 69: 699-704.
12. İlhan B, Bozdereli Berikol G, Aydın H, Arslan Erduhan M, Doğan H. COVID-19 outbreak impact on emergency trauma visits and trauma surgery in a level 3 trauma center. *Ir J Med Sci* 2022; 191: 2319-24.
13. Pintado JF, Gibaja W, Vallejos RA, Rosas W, Guerra-Farfan E, Nuñez JH. How COVID-19 has affected emergent visits to a Latin-American trauma department: Experience at a Peruvian national trauma referral center. *Injury* 2020; 51: 2834-9.
14. Rozenfeld M, Peleg K, Givon A, Bala M, Shaked G, Bahouth H, et al. COVID-19 Changed the Injury Patterns of Hospitalized Patients. *Prehosp Disaster Med* 2021; 36: 251-9.
15. Chiba H, Lewis M, Benjamin ER, Jakob DA, Liasidis P, Wong MD, et al. "Safer at home": The effect of the COVID-19 lockdown on epidemiology, resource utilization, and outcomes at a large urban trauma center. *J Trauma Acute Care Surg* 2021; 90: 708-13.
16. DiFazio LT, Curran T, Bilaniuk JW, Adams JM, Durling-Grover R, Kong K, et al. The Impact of the COVID-19 Pandemic on Hospital Admissions for Trauma and Acute Care Surgery. *Am Surg* 2020; 86: 901-3.
17. Yasin YJ, Alao DO, Grivna M, Abu-Zidan FM. Impact of the COVID-19 Pandemic on road traffic collision injury patterns and severity in Al-Ain City, United Arab Emirates. *World J Emerg Surg* 2021; 16: 57.