

The Effects of *Prunus Laurocerasus* Fruit Extract on Oxidative and Endoplasmic Reticulum Stress Responses in Doxorubicin-induced Cardiac Damage

Prunus Laurocerasus Meyve Ekstraktının Doksorubisin Tarafından İndüklenen Kardiyak Hasarda Oksidatif ve Endoplazmik Retikulum Stres Yanıtları Üzerine Etkileri

● Selma Cırrık¹, ● Emel Kabartan², ● Gülay Hacıoğlu³, ● Emine Gülçeri Güleç Peker⁴

¹Ordu University Faculty of Medicine, Department of Physiology, Ordu, Turkey

²Ordu University Ulubey Vocational School, Laboratory and Veterinary Health Program, Ordu, Turkey

³Giresun University Faculty of Medicine, Department of Physiology, Giresun, Turkey

⁴Giresun University Faculty of Engineering, Department of Basic Sciences, Giresun, Turkey

Cite this article as: Cırrık S, Kabartan E, Hacıoğlu G, Güleç Peker EG. The Effects of *Prunus Laurocerasus* Fruit Extract on Oxidative and Endoplasmic Reticulum Stress Responses in Doxorubicin-induced Cardiac Damage. J Acad Res Med

ABSTRACT

Objective: In this study, the effects of *Prunus laurocerasus* fruit extract (PLE), which is known for its high levels of phenolic compounds and antioxidant efficiency, on doxorubicin (DXR)-induced cardiotoxicity were examined.

Methods: Male Sprague-Dawley rats were randomly assigned to the Control, DXR, PLE₅₀₀+DXR, and PLE₁₀₀₀+DXR groups (n=6). PLE was orally administered to the animals in the PLE₅₀₀+DXR (500 mg/kg) and PLE₁₀₀₀+DXR (1000 mg/kg) groups for 2 weeks. DXR was injected (15 mg/kg, intraperitoneally) 2 days before sacrifice in the DXR, PLE₅₀₀+DXR, and PLE₁₀₀₀+DXR groups. At the end of the study period, serum troponin I levels, myocardial superoxide dismutase (SOD) and catalase (CAT) activities, malondialdehyde (MDA), glutathione (GSH), glucose-regulated protein (GRP)78, and pro-caspase 12 levels were measured.

Results: In the DXR group, serum troponin I concentration significantly increased (p<0.001), however PLE treatment significantly reduced this DXR-induced increment (p<0.001 and p<0.05 in the PLE₅₀₀+DXR and PLE₁₀₀₀+DXR groups, respectively). DXR-induced increase in myocardial MDA (p<0.001) significantly reduced in the PLE₅₀₀+DXR (p<0.001) and PLE₁₀₀₀+DXR (p<0.001) groups. However, DXR-induced changes, such as GSH depletion (p<0.001), reduced CAT (p<0.001) and SOD (p<0.001) activities, increased GRP78 (p<0.01) and decreased pro-caspase 12 (p<0.001) levels, were not significantly affected by PLE treatment.

Conclusion: The present results indicate that pretreatment with PLE reduced, but not completely prevented, cardiac damage induced by DXR. While a reduction in oxidative stress appears to play a role in the partial protective effect of PLE treatment, endoplasmic reticulum stress appears to have no role. Further research is necessary to understand the mechanisms of action underlying PLE treatment.

Keywords: MDA, GSH, CAT, SOD, GRP78, pro-caspase 12, troponin

ÖZ

Amaç: Bu çalışmada, yüksek düzeyde fenolik bileşik ve antioksidan etkinliğe sahip olduğu bilinen *Prunus laurocerasus* meyve özütünün (PLE), doxorubicin (DXR) tarafından indüklenen kardiyotoksisite üzerindeki etkileri incelenmiştir.

Yöntemler: Erkek Sprague-Dawley sıçanlar rastgele olarak kontrol, DXR, PLE₅₀₀+DXR ve PLE₁₀₀₀+DXR gruplarına ayrıldı (n=6). PLE₅₀₀+DXR (500 mg/kg) ve PLE₁₀₀₀+DXR (1000 mg/kg) gruplarındaki hayvanlara oral olarak iki hafta boyunca PLE uygulaması yapıldı. DXR, PLE₅₀₀+DXR ve PLE₁₀₀₀+DXR gruplarında, feda edilmeden 2 gün önce DXR enjeksiyonu yapıldı (15 mg/kg, intraperitoneal). Çalışma süresi sonunda serum troponin I düzeyleri, miyokardial superoksit dismutaz (SOD) ve katalaz (CAT) aktiviteleri ile malondialdehit (MDA), glutatyon (GSH), glukozla düzenlenen protein (GRP)78 ve pro-kaspaz 12 düzeyleri ölçüldü.

ORCID IDs of the authors: S.C. 0000-0001-8474-0185; E.K. 0000-0002-6289-2310; G.H. 0000-0002-8528-2371; E.G.G.P. 0000-0001-7244-0281.

Corresponding Author/Sorumlu Yazar: Selma Cırrık,

E-mail: selmacrrk@yahoo.com



Received Date/Geliş Tarihi: 18.11.2023 **Accepted Date/Kabul Tarihi:** 06.04.2024



Copyright© 2024 The Author. Published by Galenos Publishing House on behalf of University of Health Sciences Turkey Gaziosmanpaşa Training and Research Hospital. This is an open access article under the Creative Commons AttributionNonCommercial 4.0 International (CC BY-NC 4.0) License.

Bulgular: DXR grubunda serum troponin I konsantrasyonu önemli olarak arttı ($p<0,001$), PLE tedavisi ise DXR kaynaklı bu artışı önemli ölçüde azalttı (PLE₅₀₀+DXR ve PLE₁₀₀₀+DXR gruplarında sırasıyla $p<0,001$ ve $p<0,05$). DXR kaynaklı miyokardial MDA artışı ($p<0,001$) PLE₅₀₀+DXR ($p<0,001$) ve PLE₁₀₀₀+DXR ($p<0,001$) gruplarında önemli ölçüde azaldı. Ancak, PLE tedavisi DXR kaynaklı değişiklikleri, örneğin GSH tükenmesi ($p<0,001$), azalmış CAT ($p<0,001$) ve SOD ($p<0,001$) aktivitesi, artmış GRP78 ($p<0,01$) ve azalmış pro-kaspaz 12 ($p<0,001$) düzeylerini önemli ölçüde etkilemedi.

Sonuç: Elde edilen sonuçlar, PLE ön tedavisinin DXR tarafından indüklenen kardiyak hasarı azalttığını, ancak tamamen önlemediğini göstermektedir. PLE tedavisinin kısmi koruyucu etkisinde, oksidatif stres azalışının rol oynadığı görünse de endoplazmik retikulum stresinin rolü olmadığı anlaşılmaktadır. PLE tedavisinin temelinde yatan etki mekanizmalarını anlamak için daha fazla araştırmaya ihtiyaç vardır.

Anahtar Kelimeler: MDA, GSH, CAT, SOD, GRP78, pro-kaspaz 12, troponin

INTRODUCTION

Doxorubicin (DXR) belongs to the anthracycline group of chemotherapy drugs and is widely used for treating various cancer types, including sarcoma, lymphoma, and breast cancer. Nonetheless, its substantial efficacy is frequently overshadowed by cardiotoxic side effects, which often constrain the clinical application of DXR (1-3). Different molecular mechanisms have been suggested to underlie the development of DXR-induced cardiotoxicity, including topoisomerase II inhibition, oxidative stress, mitochondrial dysfunction, disturbances in calcium balance, and cell death (4). Among these, oxidative stress is a notable mechanism of DXR-induced cardiotoxicity. DXR increases the generation of reactive oxygen species (ROS) and weakens enzymatic and non-enzymatic antioxidant defense systems (5). Contemporary studies have demonstrated the involvement of endoplasmic reticulum (ER) stress in DXR-induced cardiotoxicity (6). ER stress is a condition arising from disruptions in ER functions due to various stressors, including adenosine triphosphate deficiency, oxidative stress, drugs, or toxins, and if it persists for a prolonged period, it can lead to cell death (7). In DXR-induced cardiotoxicity, the upregulation of ER stress proteins, including glucose-regulated protein (GRP)78, as well as caspase-12 activation have been reported previously (8,9).

To reduce the cardiotoxic effects of DXR, various strategies have been discussed, including combination therapies with different bioactive compounds (10). Phenolic compounds, including rutin, caffeic acid, chlorogenic acid, vanillic acid, quercetin, naringenin, and resveratrol, have been reported to exhibit protective effects by mitigating damage mechanisms, including oxidative stress, ER stress, or apoptosis, in DXR-induced cardiotoxicity (10-16). *Prunus laurocerasus* (*P. laurocerasus*) (synonym *Laurocerasus officinalis*), a perennial plant rich in bioactive compounds, is primarily found on the Black Sea coast of Turkey and throughout southeastern Europe and southwestern Asia (17,18). In experimental studies using *P. laurocerasus* fruit, leaf, or seed, the protective effects of extracts have been reported in conditions such as kidney and liver damage induced by paracetamol, gastric damage stimulated by indomethacin, and diabetes (17,19-23). In most cases, treatment with the extract resulted in a reduction of oxidative stress, leading to alleviation of tissue damage. Although it has the potential to be protective against DXR-induced cardiotoxicity because of its bioactive components, the possible protective effect of

P. laurocerasus against the systemic side effects of DXR has not been investigated until now. The objective of this research was to evaluate the impact of *P. laurocerasus* fruit extract (PLE) on DXR-induced cardiotoxicity through two important mechanisms in its pathophysiology: oxidative stress and ER stress responses.

METHODS

Extraction of *P. laurocerasus* Fruits

Freshly collected *P. laurocerasus* fruits were first dried and then ground into a powder. The ground fruits were mixed with 80% ethanol (1:5), and the mixture was agitated for 2 days at room temperature in a shaking water bath. The mixture was filtered through the filter paper at the end of the second day, and the fruit extract was obtained by evaporating the solvent using an evaporator at 50 °C. The concentrated extracts were diluted with tap water and administered to the treatment groups as described below.

Animals and Their Grouping

In this study, 24 male Sprague-Dawley rats weighing between 280 and 320 g were used. The Ordu University Animal Experiments Local Ethics Committee approved all the experimental procedures (decision no: 12, date: 22.10.2020). The 24 rats were randomly separated into 4 groups, namely Control, DXR, PLE₅₀₀+DXR, and PLE₁₀₀₀+DXR, with 6 rats in each group.

Control group (n=6): For 2 weeks, the animals received drinking water via intragastric gavage (ig).

DXR group (n=6): The rats were administered drinking water (ig) for 2 weeks, and DXR (15 mg/kg, single dose) was injected intraperitoneally (IP) 2 days before the end of the experimental period (24).

PLE₅₀₀+DXR group (n=6): The animals received a daily administration of PLE (500 mg/kg, ig) for 2 weeks, and DXR (15 mg/kg, IP) was administered on the 13th day of the experimental period.

PLE₁₀₀₀+DXR group (n=6): For a duration of 2 weeks, the rats orally received a daily administration of PLE (1000 mg/kg, ig), and DXR (15 mg/kg, IP) was injected on the 13th day of the study.

On the 15th day of the study, the rats were anesthetized with ketamine and xylazine (80 mg/kg: 10 mg/kg, IP). Blood samples were collected from the abdominal aorta and allowed to

coagulate at ambient temperature. Subsequently, serum samples were obtained by centrifugation at 1000 g for 20 min. Heart tissue and serum samples were kept in a 80 °C deep freezer until these parameters were analyzed:

Serum troponin I: The level of cardiac troponin I in the serum samples was measured using a commercial enzyme-linked immunosorbent assay kit (Elabscience, E-EL-R1253, China). The analysis was performed in accordance with the guidelines provided by the manufacturer, and the outcomes were evaluated by comparing the optical density of the samples against the standard curve.

Malondialdehyde (MDA): Myocardial MDA levels were determined using Buege and Aust's (25) thiobarbituric acid reactive substance formation method. The pink color formed by the reaction of MDA with thiobarbituric acid was detected at 535 nm using a spectrophotometer (25).

Glutathione (GSH): GSH levels in cardiac homogenates were measured by the method of Aykaç et al. (26). The yellow color formed by the interaction of GSH and 5,5-dithio-bis (2-nitrobenzoic acid) was detected using a spectrophotometer at 412 nm.

Catalase (CAT) activity: CAT activity in cardiac tissue was assessed by measuring absorbance at 240 nm, reflecting the rate of hydrogen peroxide (H₂O₂) decomposition catalyzed by CAT per unit of time (27).

Superoxide dismutase (SOD) activity: SOD activity in cardiac tissue was measured by determining the red color created by superoxide radicals, which are formed using xanthine and xanthine oxidase by reducing nitroblue tetrazolium at 505 nm (28).

GRP78: GRP78 levels were measured using commercial ELISA kits (Bioassay Technology Laboratory BT-E1255Ra, China) in cardiac tissue homogenates. The assay protocol was performed in accordance with the manufacturer's guidelines. The results were computed using a standard curve and conveyed as ng/mg of protein.

Pro-caspase 12: Myocardial pro-caspase 12 concentrations were assayed using a commercial ELISA kit (Bioassay Technology Laboratory BT-E2111Ra, China) according to the manufacturer's instructions. The pro-caspase 12 concentration in tissue homogenates was determined by contrasting the optical density of the samples with the standard curve, and the findings were expressed in relation to the total protein content.

Statistical Analysis

The findings are presented as mean ± standard deviation. Normality and homogeneity of the data were assessed using the Kolmogorov-Smirnov and Levene's tests, respectively. Normally distributed and homogeneous data were then tested by One-Way analysis of variance, followed by the Tukey test as a post hoc analysis. Significance was assigned to values with p<0.05

RESULTS

Serum troponin I: The serum troponin I level was 806.75±52.72 pg/mL in the control group. This value was significantly higher

in the DXR (1813.68±142.36 pg/mL, p<0.001), PLE₅₀₀+DXR (1554.34±174.63 pg/mL, p<0.001) and PLE₁₀₀₀+DXR (1514.03±141.74 pg/mL, p<0.001) groups. Compared with the DXR group, PLE₅₀₀+DXR (p<0.05) and PLE₁₀₀₀+DXR (p<0.01) groups had higher levels of troponin I. No statistical difference was observed between the two extract treatment groups (Figure 1).

Oxidative Stress Parameters

In the control group, the MDA level was measured as 188.28±45.85 nmol/g wet tissue (wt). Increased concentrations of MDA were observed in the DXR (527.67±64.68 nmol/g wt, p<0.001), PLE₅₀₀+DXR (343.245±47.99 nmol/g wt, p<0.001) and PLE₁₀₀₀+DXR (368.71±72.26 nmol/g wt, p<0.001) groups. In the PLE₅₀₀+DXR and PLE₁₀₀₀+DXR groups, MDA levels were significantly lower than those in the DXR group (p<0.001 and p<0.001, respectively) (Figure 2A).

GSH levels were significantly lower in the DXR (1.13±0.26 µmol/g wt, p<0.001), PLE₅₀₀+DXR (1.42±0.32 µmol/g wt, p<0.001) and PLE₁₀₀₀+DXR (1.25±0.34 µmol/g wt, p<0.001) groups than in the control values (3.82±0.59 µmol/g wt). Treatment with two different doses of PLE did not significantly change cardiac GSH levels, and the results were not different from the DXR group or each other (Figure 2B).

Cardiac CAT activity was 3.7±0.36 µmol/H₂O₂/min/mg wt in the control group. The value significantly reduced to 1.412±0.41 µmol/H₂O₂/min/mg wt in the DXR group (p<0.001). CAT activities were significantly lower in the PLE₅₀₀+DXR (1.79±0.38 µmol/H₂O₂/min/mg wt) and PLE₁₀₀₀+DXR (1.62±0.47 µmol/H₂O₂/min/mg wt) groups compared with the control (p<0.001 and p<0.001, respectively), and the values were not significantly different from the DXR group or each other (Figure 2C).

SOD activity was measured as 4.24±0.53 U/mg wt in the control group and was significantly reduced in the DXR (1.37±0.50 U/mg wt, p<0.001), PLE₅₀₀+DXR (1.64±0.57 U/mg wt, p<0.001)




Figure 1. Circulating troponin I concentrations in the control, DXR, PLE₅₀₀+DXR and PLE₁₀₀₀+DXR groups (n=6). The results were presented as the mean ± standard deviation. Comparing to the control group *p<0.001, comparing to the DXR group #p<0.05, ##p<0.01.

PLE: *Prunus laurocerasus* fruit extract, DXR: doxorubicin

and PLE₁₀₀₀+DXR (1.82±0.539 /mg wt, p<0.001) groups. Compared with the DXR group, no significant alterations were noted in the PLE₅₀₀+DXR and PLE₁₀₀₀+DXR groups (Figure 2D).

ER Stress Parameters

The GRP78 level was 0.59±0.07 ng/mg prot in the control group and significantly increased to 0.79±0.08 ng/mg prot in the DXR group (p<0.01). Although the GRP78 levels were lower in the PLE₅₀₀+DXR (0.68±0.13 ng/mg prot) and PLE₁₀₀₀+DXR (0.64±0.11 ng/mg prot) groups, the results were not significantly different from the control or DXR groups (Figure 3A).

The pro-caspase 12 level in the DXR-injected group (10.14±1.57 ng/mg prot) was found to be significantly lower (p<0.01),

than that in the control group (16.32±3.43 ng/mg prot). In the PLE₅₀₀+DXR and PLE₁₀₀₀+DXR groups, pro-caspase 12 levels did not significantly change (12.91±1.82 and 13.19±2.93 ng/mg prot, respectively) (Figure 3B).

DISCUSSION

Cancer remains a major global health challenge, with a growing number of individuals being diagnosed due to rising life expectancy and population growth. Therefore, research is ongoing to reduce the cardiotoxic side effects of DXR, a widely used anticancer drug. This study investigated the effectiveness of PLE in reducing the cardiotoxic side effects of DXR.




Figure 2. Oxidative stress parameters measured in myocardial tissue of the control, DXR, PLE₅₀₀+DXR and PLE₁₀₀₀+DXR groups (n=6). Myocardial MDA levels (A), GSH levels (B), CAT activity (C) and SOD activity (D). The results were presented as the mean ± standard deviation. Comparing to the control group *p<0.001, comparing to the DXR group #p<0.001. PLE: *Prunus laurocerasus* fruit extract, DXR: doxorubicin, CAT: catalase, GSH: glutathione, MDA: malondialdehyde, SOD: superoxide dismutase




Figure 3. ER stress parameters measured in myocardial tissue of the control, DXR, PLE₅₀₀+DXR and PLE₁₀₀₀+DXR groups (n=6). Myocardial GRP78 levels (A), pro-caspase 12 levels (B). Comparing to the control group *p<0.01. PLE: *Prunus laurocerasus* fruit extract, DXR: doxorubicin, GRP78: glucose-regulated protein78

Myocardial damage induced by DXR can be monitored using various circulating biomarkers, including cardiac troponins. The cardiac-specific troponin proteins (troponin I, troponin T) are located in the myocardium, enabling their use as specific biomarkers to identify alterations in cardiac muscle. Because troponin I is accepted as a sensitive and specific indicator of DXR-induced myocardial injury, its serum concentrations were evaluated in the present study (29,30). Increased troponin I levels in the DXR group were considered an indicator of cardiac damage. Oxidative stress is a conspicuous mechanism in the pathophysiology of DXR-induced cardiotoxicity (5,31). Elevated ROS levels lead to modifications in the functional and structural components of the cell, resulting in myocardial damage and functional loss. Previous studies have reported that a decrease in cardiac antioxidant enzymes such as CAT, SOD, and GSH peroxidase and a reduction in the level of the non-enzymatic antioxidant molecule GSH are important factors in the development of oxidative damage of the myocardium during DXR treatment (5,31,32). In the current study, the increase in lipid peroxidation, along with GSH depletion and decreased SOD and CAT activities, suggested the presence of DXR-induced cardiac oxidative stress.

The *P. laurocerasus* fruit used in this study is rich in phenolic compounds such as chlorogenic, caffeic, gallic, vanillic, benzoic, and coumaric acids, rutin, and quercetin. This high phenolic content largely contributes to the pharmaceutical significance of the plant (33,34). In studies using different experimental damage models, it has been shown that pretreatment with *P. laurocerasus* extracts successfully reduces injury-related oxidative stress in stomach, kidney, and liver tissues (17-22). Not only fruit extract but also seed and leaf extracts of *P. laurocerasus* have been used in previous studies. For example, Uslu and Uslu (23) demonstrated a significant improvement in oxidative stress parameters (MDA, GSH peroxidase, GSH) in diabetic rats treated with leaf, fruit, or seed extracts of *P. laurocerasus* (21 days, 500 mg/kg) and reported that the effectiveness of these different extracts was at the similar level. To date, it has not been shown whether *P. laurocerasus* extracts affect cardiac oxidative stress in any injury model. Our study is the first to demonstrate that treatment with fruit extracts of *P. laurocerasus* can reduce oxidative stress in cardiac tissue. According to the present results, lipid peroxidation induced by DXR significantly decreased in the extract treatment groups; however, it did not return to control values. Similar changes observed in serum troponin levels indicate that the treatment could not completely prevent oxidative damage of the myocardium but was successful in reducing it. Decreased lipid peroxidation after *P. laurocerasus* treatment appeared to be related to the free radical scavenging properties of the extract because there was no improvement in endogenous GSH levels or antioxidant enzyme activities (CAT, SOD). This finding contradicts previous studies demonstrating elevated GSH levels and increased SOD and CAT activities in renal, gastric, and hepatic tissues following *P. laurocerasus* treatment (17-22), indicating that the efficacy of *P. laurocerasus* might vary depending on the tissue, as increasing

the dose of the extract did not alter its efficiency for the heart.

The ER plays a crucial role in eukaryotic cells, being responsible for tasks such as the synthesis, folding, and secretion of proteins and maintaining calcium homeostasis. ER stress is a cellular response aimed at restoring ER functions and generally involves a slowing down of protein synthesis, accelerated proteasomal degradation of unfolded or misfolded proteins, and increased synthesis of ER chaperone proteins involved in the protein folding process, such as GRP78 (7). In the current study, the increased GRP78 level in the DXR group indicated the presence of cardiac ER stress, which is consistent with previous studies demonstrating the role of ER stress in DXR-induced cardiotoxicity (8,9,35). Although restoration of cellular homeostasis is aimed, severe or prolonged ER stress triggers cell death through apoptotic signaling pathways, namely caspase-12, c-JUN NH2-terminal kinase, and C/EBP homologous protein pathways. In the present study, a significant decrease in pro-caspase 12 levels was interpreted as an increase in caspase 12 activation, a result that is consistent with previous literature findings indicating the role of ER stress-mediated apoptosis in DXR-induced cardiotoxicity (8,9,35). While the effect of PLE on oxidative stress has been previously studied for different tissues, excluding the heart, its effect on ER stress in the heart was examined for the first time in this study. Our results showed a trend for GRP78 and pro-caspase 12 levels to return to the control levels, but these changes caused by extract pretreatment were not statistically significant in the DXR group. Although individual bioactive components within *P. laurocerasus* have been reported to reduce DXR-induced ER stress (10-16), it appears that their levels within the extract may not be sufficient to produce the same response even at higher doses. Because oxidative stress is one of the ER stress triggering factors, reduced oxidative stress would be expected to be accompanied by decreased ER stress in the extract-treated groups. However, it should be noted that oxidative stress was not completely prevented in the extract groups. Ongoing oxidative stress may play a significant role in the persistence of both ER stress and cardiac damage.

Study Limitations

One of the most significant shortcomings of this study is that cardiac damage was assessed only through serum troponin I measurement without histopathological evaluation. Demonstrating the histological evidence of both DOX-induced damage and partial recovery after treatment could have supported the current findings.

CONCLUSION

In this study, the effects of PLE on DOX-induced cardiotoxicity were investigated for the first time. Although histopathological evaluation was not conducted, which is a limitation of the current study, the results indicated that pretreatment with PLE partially prevented myocardial damage in DXR-induced cardiotoxicity. We suggested that PLE treatment decreased cardiac lipid peroxidation, possibly due to its endogenous radical scavenging

property, without causing any changes in the tissue levels of enzymatic (SOD and CAT) or non-enzymatic antioxidants (GSH). While a reduction in oxidative stress appears to play a role in the partial protection of PLE treatment, it seems that the ER stress response plays no role. The incomplete prevention of DXR-induced cardiac damage does not appear to be related to dose insufficiency; however, future studies with longer treatment durations can be planned.

Ethics Committee Approval: The Ordu University Animal Experiments Local Ethics Committee approved all the experimental procedures (decision no: 12, date: 22.10.2020).

Informed Consent: Animal experiment study.

Author Contributions: Surgical and Medical Practices - S.C., E.K., G.H.; Concept - S.C.; Design - S.C.; Data Collection and/or Processing - S.C., E.K., G.H., E.G.G.P.; Analysis and/or Interpretation - S.C., E.K., G.H., E.G.G.P.; Literature Search - S.C., E.K.; Writing - S.C.

Conflict of Interest: The authors have no conflict of interest to declare.

Financial Disclosure: Financial assistance for this study was provided by Ordu University (Grant number A2101).

Etik Komite Onayı: Ordu Üniversitesi Hayvan Deneyleri Yerel Etik Kurulu tüm deney prosedürlerini onayladı (karar no: 12, tarih: 22.10.2020).

Hasta Onamı: Hayvan deneyi çalışması.

Yazar Katkıları: Cerrahi ve Medikal Uygulama - S.C., E.K., G.H.; Konsept - S.C.; Dizayn - S.C.; Veri Toplama veya İşleme - S.C., E.K., G.H., E.G.G.P.; Analiz veya Yorumlama - S.C., E.K., G.H., E.G.G.P.; Literatür Arama - S.C., E.K.; Yazan - S.C.

Çıkar Çatışması: Yazarlar tarafından çıkar çatışması bildirilmemiştir.

Finansal Destek: Bu çalışma için finansal destek Ordu Üniversitesi tarafından sağlanmıştır (hibe numarası A2101).

REFERENCES

1. Wu BB, Leung KT, Poon EN. Mitochondrial-Targeted Therapy for Doxorubicin-Induced Cardiotoxicity. *Int J Mol Sci* 2022; 23: 1912.
2. Christidi E, Brunham LR. Regulated cell death pathways in doxorubicin-induced cardiotoxicity. *Cell Death Dis* 2021; 12: 339.
3. Varela-López A, Battino M, Navarro-Hortal MD, Giampieri F, Forbes-Hernández TY, Romero-Márquez JM, et al. An update on the mechanisms related to cell death and toxicity of doxorubicin and the protective role of nutrients. *Food Chem Toxicol* 2019; 134: 110834.
4. Rawat PS, Jaiswal A, Khurana A, Bhatti JS, Navik U. Doxorubicin-induced cardiotoxicity: An update on the molecular mechanism and novel therapeutic strategies for effective management. *Biomed Pharmacother* 2021; 139: 111708.
5. Songbo M, Lang H, Xinyong C, Bin X, Ping Z, Liang S. Oxidative stress injury in doxorubicin-induced cardiotoxicity. *Toxicol Lett* 2019; 307: 41-8.
6. Yarmohammadi F, Rezaee R, Haye AW, Karimi G. Endoplasmic reticulum stress in doxorubicin-induced cardiotoxicity may be therapeutically targeted by natural and chemical compounds: A review. *Pharmacol Res* 2021; 164: 105383.
7. Tabas I, Ron D. Integrating the mechanisms of apoptosis induced by endoplasmic reticulum stress. *Nat Cell Biol* 2011; 13: 184-90.
8. Kim BS, Park IH, Lee AH, Kim HJ, Lim YH, Shin JH. Sacubitril/valsartan reduces endoplasmic reticulum stress in a rat model of doxorubicin-induced cardiotoxicity. *Arch Toxicol* 2022; 96: 1065-74.
9. Wang Z, Wang M, Liu J, Ye J, Jiang H, Xu Y, et al. Inhibition of TRPA1 attenuates doxorubicin-induced acute cardiotoxicity by suppressing oxidative stress, the inflammatory response, and endoplasmic reticulum stress. *Oxid Med Cell Longev* 2018; 2018: 5179468.
10. Syahputra RA, Harahap U, Dalimunthe A, Nasution MP, Satria D. The Role of Flavonoids as a Cardioprotective Strategy against Doxorubicin-Induced Cardiotoxicity: A Review. *Molecules* 2022; 27: 1320.
11. Cicek B, Hacimuftuoglu A, Yeni Y, Danisman B, Ozkaraca M, Mokhtare B, et al. Chlorogenic acid attenuates doxorubicin-induced oxidative stress and markers of apoptosis in cardiomyocytes via Nrf2/HO-1 and dityrosine signaling. *J Pers Med* 2023; 13: 649.
12. Zhang Y, Kong D, Han H, Cao Y, Zhu H, Cui G. Caffeic acid phenethyl ester protects against doxorubicin-induced cardiotoxicity and increases chemotherapeutic efficacy by regulating the unfolded protein response. *Food Chem Toxicol* 2022; 159: 112770.
13. Hu LF, Lan HR, Li XM, Jin KT. A Systematic Review of the Potential Chemoprotective Effects of Resveratrol on Doxorubicin-Induced Cardiotoxicity: Focus on the Antioxidant, Antiapoptotic, and Anti-Inflammatory Activities. *Oxid Med Cell Longev* 2021; 2021: 2951697.
14. Baniahmad B, Safaeian L, Vaseghi G, Rabbani M, Mohammadi B. Cardioprotective effect of vanillic acid against doxorubicin-induced cardiotoxicity in rat. *Res Pharm Sci* 2020; 15: 87-96.
15. Ma Y, Yang L, Ma J, Lu L, Wang X, Ren J, et al. Rutin attenuates doxorubicin-induced cardiotoxicity via regulating autophagy and apoptosis. *Biochim Biophys Acta Mol Basis Dis* 2017; 1863: 1904-11.
16. Dong Q, Chen L, Lu Q, Sharma S, Li L, Morimoto S, et al. Quercetin attenuates doxorubicin cardiotoxicity by modulating Bmi-1 expression. *Br J Pharmacol* 2014; 171: 4440-54.
17. Uslu H, Atila G, Özen H, Karaman M. Effects of different doses of Prunus laurocerasus L. leaf extract on oxidative stress, hyperglycaemia and hyperlipidaemia induced by type I diabetes. *Indian J Tradit Knowl* 2018; 17: 430-6.
18. Elmastas M, Genc N, Demirtas I, Aksit H, Aboul-Enien HY. Isolation and identification of functional components in seed of cherry laurel (*Laurocerasus officinalis* Roem.) and investigation of their antioxidant capacity. *J Biologically Act Prod Nat* 2013; 3: 115-20.
19. Turan MI, Turkoglu M, Dundar C, Celik N, Suleyman H. Investigating the effect of Prunus laurocerasus fruit extract in type II diabetes induced rats. *Int J Pharmacol* 2013; 9: 373-8.
20. Aydin Bertkas O, Keskin A, Gulec Peker EG. Gastroprotective effects of Prunus laurocerasus L. fruit extracts against the oxidative stress induced by indomethacin in rats. *Eur J Clin Exp Med* 2022; 3: 284-9.
21. Kaltalioglu K. Prunus laurocerasus L. extracts prevent paracetamol induced nephrotoxicity by regulating antioxidant status: An experimental animal model. *Hittite Journal of Science and Engineering* 2022; 9: 275-80.
22. Bertkas OA, Peker EGG. Protective effects of Prunus laurocerasus extracts against paracetamol-induced hepatotoxicity. *J Nutrition and Food Processing* 2022; 5: 1-5.
23. Uslu H, Uslu GA. Evaluating the effects of Prunus laurocerasus seed, fruit and leaf extracts on hyperglycaemia, insulin sensitivity and anti-oxidative activities in experimental diabetes in rat. *Thai J Vet Med* 2021; 51: 667-73.
24. Kelleni MT, Amin EF, Abdelrahman AM. Effect of Metformin and Sitagliptin on Doxorubicin-Induced Cardiotoxicity in Rats: Impact of Oxidative Stress, Inflammation, and Apoptosis. *J Toxicol* 2015; 2015: 424813.
25. Buege JA, Aust SD. Microsomal lipid peroxidation. *Methods Enzymol* 1978; 52: 302-10.
26. Aykaç G, Uysal M, Yalçın AS, Koçak-Toker N, Sivas A, Oz H. The effect of chronic ethanol ingestion on hepatic lipid peroxide, glutathione, glutathione peroxidase and glutathione transferase in rats. *Toxicology* 1985; 36: 71-6.
27. Aebi H. Catalase in vitro. *Methods Enzymol* 1984; 105: 121-6.
28. Sun Y, Oberley LW, Li Y. A simple method for clinical assay of superoxide dismutase. *Clin Chem* 1988; 34: 497-500.
29. Upadhyay S, Gupta KB, Mantha AK, Dhiman M. A short review: Doxorubicin and its effect on cardiac proteins. *J Cell Biochem* 2021; 122: 153-65.
30. Atas E, Kismet E, Kesik V, Karaoglu B, Aydemir G, Korkmazer N, et al. Cardiac troponin-I, brain natriuretic peptide and endothelin-1 levels in a rat model of doxorubicin-induced cardiac injury. *J Cancer Res Ther* 2015; 11: 882-6.
31. Zhang YY, Yi M, Huang YP. Oxymatrine Ameliorates Doxorubicin-Induced Cardiotoxicity in Rats. *Cell Physiol Biochem* 2017; 43: 626-35.
32. Wu YZ, Zhang L, Wu ZX, Shan TT, Xiong C. Berberine Ameliorates Doxorubicin-Induced Cardiotoxicity via a SIRT1/p66Shc-Mediated Pathway. *Oxid Med Cell Longev* 2019; 2019: 2150394.
33. Ozturk B, Celik SM, Karakaya M, Karakaya O, Islam A, Yarılgac T. Storage temperature affects phenolic content, antioxidant activity and fruit quality parameters of cherry laurel (*Prunus laurocerasus* L.). *Journal of Food Processing and Preservation* 2017; 41: e12774.

34. Karabegović IT, Stojičević SS, Veličković DT, Todorović ZB, Nikolić NČ, Lazić ML. The effect of different extraction techniques on the composition and antioxidant activity of cherry laurel (*Prunus laurocerasus*) leaf and fruit extracts. *Industrial Crops and Products* 2014; 54: 142-8.
35. Lou Y, Wang Z, Xu Y, Zhou P, Cao J, Li Y, et al. Resveratrol prevents doxorubicin-induced cardiotoxicity in H9c2 cells through the inhibition of endoplasmic reticulum stress and the activation of the Sirt1 pathway. *Int J Mol Med* 2015; 36: 873-80.